

Reconsidering the origin of the 21 micron feature: Oxides in carbon-rich PPNe? ¹

Th. Posch

Institut für Astronomie, Türkenschanzstraße 17, A-1180 Wien, Austria

posch@astro.univie.ac.at

H. Mutschke

Astrophysikalisches Institut, Schillergässchen 2-3, D-07745 Jena, Germany

and

A. Andersen

NORDITA, Blegdamsvej 17, DK-2100 Copenhagen, Denmark

ABSTRACT

The origin of the so-called ‘21’ μm feature which is especially prominent in the spectra of some carbon-rich protoplanetary nebulae (PPNe) is the matter of a lively debate. A large number of potential band carriers have been presented and discarded within the past decade. The present paper gives an overview of the problems related to the hitherto proposed feature identifications, including the recently suggested candidate carrier silicon carbide. We also discuss the case for spectroscopically promising oxides.

SiC is shown to produce a strong resonance band at 20–21 μm if coated by a layer of silicon dioxide. At low temperatures, core-mantle particles composed of SiC and amorphous SiO₂ indeed have their strongest spectral signature at a position of 20.1 μm , which coincides with the position of the ‘21’ μm emission band.

The optical constants of another candidate carrier that has been relatively neglected so far – iron monoxide – are proven to permit a fairly accurate reproduction of the ‘21’ μm feature profile as well, especially when low-temperature measurements of the infrared properties of FeO are taken into account. As candidate carrier of the ‘21’ μm emission band, FeO has the advantage of being stable against further oxidation and reduction only in a narrow range of chemical and physical conditions, coinciding with the fact that the feature, too, is detected in a small group of objects only. However, it is unclear how FeO should form or survive particularly in carbon-rich PPNe.

Subject headings: circumstellar matter — infrared: stars — methods: laboratory — stars: AGB and post-AGB — stars: atmospheres

1. Introduction

Kwok, Volk & Hrivnak (1989) reported the discovery of a strong emission band, centered (as it then seemed) at $21\,\mu\text{m}$, in the spectra of four carbon-rich protoplanetary nebulae (PPNe) taken by the Infrared Astronomical Satellite (IRAS). The feature was not resolved well with the Low Resolution Spectrometer (LRS), but nevertheless, it was evident already from the LRS observations that it has a considerable strength in some objects (especially IRAS 07134+1005) and is too broad (full width at half maximum $>2\,\mu\text{m}$) to originate from any atomic transition. The pioneering work by Kwok, Volk & Hrivnak also highlighted the fact that the ‘21’ μm feature only occurs in a short transitional phase of stellar evolution, i.e. in a very limited range of physical and chemical conditions. Any identification of the ‘21’ μm feature has to cope with this fact.

Observations of selected PPNe with the Infrared Space Observatory (ISO) by Hrivnak, Volk & Kwok (2000) brought substantial progress, especially concerning the position and width of the feature, which was shown to be $20.1\,\mu\text{m}$ instead of $21\,\mu\text{m}$ (notwithstanding, the feature is still being referred to as ‘21’ μm band, a tradition to which we also stick in this paper). Hrivnak, Volk & Kwok (2000) analysed ISO observations at $2\text{--}45\,\mu\text{m}$ for seven PPNe and two other carbon-rich objects. Six of the sources exhibit emission features at

21 and $30\,\mu\text{m}$. The authors resolved the $30\,\mu\text{m}$ band into a broad feature peaking at $27.2\,\mu\text{m}$ (the ‘30’ μm feature) and a narrower feature at $25.5\,\mu\text{m}$ (the ‘26’ μm feature). The observations suggest that the sources with ‘21’ μm feature do also show the $30\,\mu\text{m}$ feature. The ‘26’ μm feature appears to be present in a broader range of objects – carbon stars, PPNe, and possibly PNe – than does the ‘21’ μm feature.

The candidate carriers of the ‘21’ μm band have since its discovery been the matter of a long discussion. Cox (1990) presented IRAS-LRS-spectra of ten HII regions which showed a peak close to $21\,\mu\text{m}$. He associated this peak with the feature discovered by Kwok, Volk & Hrivnak and tentatively assigned it to iron oxides with the stoichiometries $\gamma\text{-Fe}_2\text{O}_3$ and Fe_3O_4 . In the late 1990s, some of these HII regions have also been observed with ISO; in the ISO spectra, however, we do not find any traces of a broad 21 or $20\,\mu\text{m}$ emission band (in accordance with Oudmaijer & de Winter (1995), who already cast doubt on Cox’ observations).

Goebel (1993), still on the basis of IRAS spectra, assigned the ‘21’ μm feature to silicon disulfide (SiS_2), which has indeed an intense vibrational band in the $20\,\mu\text{m}$ spectral range (but, unfortunately, a second one which is not observed; see below, Sect. 4). From a chemical point of view, other sulfur compounds like S_8 , OCS , S_2 and CS_2 , even though mainly as mantles of other grains, were also taken into account by Goebel as potential emitters in the $20\,\mu\text{m}$ region.

Webster (1995) brought fullerenes as potential carriers of the ‘21’ μm feature into the discussion. Hill, Jones & d’Hende-

¹Based on observations with ISO, an ESA project with instruments funded by ESA Member States (especially the PI countries: France, Germany, the Netherlands and the United Kingdom) and with the participation of ISAS and NASA.

court (1998), having studied the IR bands of terrestrial nitrogen-doped and neutron-irradiated diamonds, showed that defects in diamond structures can cause an emission band at 20–22 μm and even stronger bands in the 8–10 μm region. However, there is a significant discrepancy between the position of 21–22 μm and the actual position of the ‘21’ μm band as determined by recent observations.

Papoular (2000) calculated synthetic spectra of carbonaceous macromolecules containing OH, oxygen, sulfur, nitrogen or a combination of them and found that it is possible to reproduce not only the ‘21’ μm feature, but also the 3.3, 11.3, 12+, 26 and 30 μm bands by emissions of such molecules. According to this scenario, the ‘21’ μm feature would arise from out-of-plane vibrations of oxygen or nitrogen atoms in 5-membered carbon rings. Papoular’s approach seems promising because it aims at an identification of *all* the major features detected in the sources of the ‘21’ μm band on the basis of *one* unified dust model. But at the time being, substantial problems remain, such as the unsatisfactory reproduction of the observed 26–35 μm band profile.

There are various other papers discussing carbonaceous materials as carriers of the ‘21’ μm feature, but without any detailed fits of its average profile. Sourisseau, Coddens & Papoular (1992) suggested amides, especially urea, as potential carrier, Buss et al. (1990) and Justtanont et al. (1996) suggested polycyclic aromatic hydrocarbon molecules and Grishko et al. (2001) hydrogenated amorphous carbon.

Von Helden et al. (2000) came up with another suggestion, attributing the

20.1 μm feature to titanium carbide clusters (made of 27 to 125 atoms). Spectra of bulk titanium carbide and TiC particles as those subsequently published by Henning & Mutschke (2001) and Kimura & Kaito (2003) show no band in the respective wavelength region, but rather weak bands at 9.5, 12.5 (Kimura & Kaito 2003) and 19 μm (Henning & Mutschke 2001; see also Speck & Hofmeister 2004). This does not contradict the measurements by von Helden et al. (2000), however, since clusters are likely to have electronic and vibrational properties deviating from those of bulk material even if composition and crystal structure are similar.

Notwithstanding, in 2003, three papers doubting the ‘TiC-cluster-identification’ with substantial arguments came out, all based essentially on arguments concerning the abundance of titanium (Hony et al. 2003, Chigai et al. 2003, and Li 2003). Hony et al. (2003) argue that TiC, like any kind of dust, can only emit as much in the infrared as it absorbs in the UV and visual range. Assigning a simple model opacity to TiC at short wavelengths, Hony et al. found that this energy balance can be fulfilled only if unrealistically large amounts of TiC are predicted to form in carbon-rich PPNe. This point was strengthened by Li (2003) by applying a sum-rule based on Kramers-Kronig considerations on the wavelength-integrated absorptivity. Chigai et al. (2003) came to the same result, comparing the number densities of SiC and TiC grains necessary for producing the 11 μm and 21 μm features, respectively, with the available Si and Ti abundances.

Recently, Speck & Hofmeister (2004) reported the experimental finding that sili-

con carbide, under certain circumstances, does not only show the well-known resonance feature at $11\,\mu\text{m}$, but also a secondary band (not predicted by group theory), which is centered at $20\text{--}21\,\mu\text{m}$. This secondary band is reported to appear only in the β -SiC-polytype, and nitrogen or carbon impurities were suspected by the authors to favour its occurrence. Cold, maybe N- or C-doped β -SiC-dust (with temperatures around 100 K) would then, according to Speck & Hofmeister, emit at $20\text{--}21\,\mu\text{m}$ (depending on particle size and shape) even more effectively than at $11\,\mu\text{m}$, since the latter peak would still be in the Wien regime of the corresponding Planck function. Important as this suggestion might be for the identification of the ‘ $21\,\mu\text{m}$ band’, it is not beyond any doubt, as we try to show in §3.

The present paper is structured as follows. In §2, we summarize the currently available astronomical information on the sources in which the enigmatic ‘ $21\,\mu\text{m}$ band’ is observed. §3 presents new insights to the infrared properties of materials which should, despite some problems, *not* be excluded as feature carriers – such as core-mantle particles composed of SiC and SiO_2 and cold FeO dust grains. In §4, we deliver additional arguments for ruling out potential carriers of this band which have been ruled out more or less convincingly by other authors. The last section (§5) summarizes our conclusions.

2. Properties of the sources of the ‘ $21\,\mu\text{m}$ feature’

So far, the ‘ $21\,\mu\text{m}$ feature’ is observed only in emission, only in objects with a C/O-ratio close to or greater than unity

and only in *cool* dusty environments, i.e. environments with dust temperatures below 250 K. Among them are the already mentioned C-rich PPNe (about half of them show the feature), two Planetary Nebulae (Hony, Waters & Tielens 2001) with central stars of the Wolf-Rayet type and, according to Volk, Xiong & Kwok (2000), two extreme carbon stars (IRAS 21318+5631, IRAS 06582+1507). Clément et al. (2004), however, re-analyzing the ISO-spectrum of IRAS 21318+5631, find no evidence for a feature at $20\,\mu\text{m}$, and in the case of IRAS 06582+1507, Volk, Xiong & Kwok (2000) also remark that its reality is doubtful.

It should be noted that an emission band with a position and width quite comparable to the ‘ $21\,\mu\text{m}$ band’ is detected in the spectra of stars on the *early* Asymptotic Giant Branch (AGB), i.e. at the very opposite end of AGB evolution: the $19.5\,\mu\text{m}$ feature (Posch et al. 2002). Its peak position amounts to $19.4\text{--}19.6\,\mu\text{m}$, its profile is symmetric and its full width at half maximum (FWHM) is close to $3\,\mu\text{m}$. The dust shells of the mostly semiregular variable stars in which this feature is observed are generally assumed to consist of refractory oxides, but not at all of carbonaceous dust grains. A solid solution of FeO and MgO has been proposed as feature carrier in this case (see also Ferrarotti & Gail (2003)).

Coming back to the PPNe exhibiting the $21\,\mu\text{m}$ band in its ‘canonical’ form, let us consider the spectrum and the derived properties of its best-known source, IRAS 07134+1005 (=SAO 96709 or HD 56126, see Fig. 1). The F5 I central star of this PPN has a surface temperature of about




Fig. 1.— ISO-SWS spectrum of the PPN IRAS 07134+1005 at a resolution $\lambda/\Delta\lambda = 600$. At the bottom, the profile of the ‘21’ μm band is shown, both as derived from the ISO-SWS spectrum HD 56126 (solid line) and from an ISO-SWS spectrum of SAO 34504 (open triangles) by subtraction of a polynomial (dotted line overplotted to the ISO spectrum at the top).

7000 K according to Hrivnak, Volk & Kwok (2000); other PPNe sources of the same feature have comparable effective temperatures (5500 to 8000 K). The circumstellar envelopes of these objects are all *optically thin* at infrared wavelengths. As for unusual properties of the central stars, van Winckel & Reyniers (2000) found them to be enriched in s-process elements and metal-poor. Apart from the peak due to the radiation of the central star at visual wavelengths, several infrared bands arising from dust grains are present in the spectrum of HD 56126, namely at 8, 11.3, 12.1, and 30 μm . They are partly unidentified, partly attributed to SiC (11.3–broad component), PAHs (11.3–narrow component, 12+) and magnesium sulfide (30 μm), respectively. No silicate features are seen in any of the sources of the ‘21’ μm band.

We derived the profile of the ‘21’ μm band using the polynomial subtraction method which has been applied by Posch et al. (1999) for the derivation of the 13 μm band profile. The residual emission of HD 56126, based on an ISO-SWS-spectrum reduced with the ISO Spectral Analysis Package (ISAP), is shown at the bottom of Fig. 1. For comparison, the scaled residual emission of SAO 34504 – calculated by the same method – is shown in the bottom part of Fig. 1 as well (open triangles). The close coincidence of the two emission profiles is noteworthy; together with the result by Volk, Kwok & Hrivnak (1999) that the feature profile is pretty much the same in all sources, this implies that the spectrum of HD 56126 can be considered representative and will be so considered in the following. The most conspicuous properties of the feature profile are a steep rise in its blue wing, a maximum position of 20.1 μm and a long ‘tail’ extending to 23–24 μm .

In contrast to stars on the early AGB, bright PPNe have spatial extensions which are large enough to resolve them even at infrared wavelengths. Kwok, Volk & Hrivnak (2002) and Miyata et al. (2003) tried to answer the question whether the different broad infrared features of PPNe have their origin in different or coincident regions on the basis of imaging through broad filters. They came to the conclusion that the 11.3 μm SiC, the 12+ μm plateau emission, and the ‘21’ μm emission originate (approximately) in the same region, which seems to indicate that similar physical and chemical conditions are required for the formation and/or survival of the respective feature carrier.

3. Silicon carbide, silicon dioxide and iron monoxide

3.1. Silicon carbide with impurities or coatings

As mentioned in §1, Speck & Hofmeister (2004) suggested silicon carbide with impurities as carrier both of the 11.3 and the ‘21’ μm spectral features. One of their arguments in favour of this hypothesis is the spatial co-location of both emission bands in some of the objects exhibiting them. Obviously, this argument is applicable not only to N-doped and C-doped SiC, but also to any other particular SiC grain population like e.g. oxidized SiC grains (see below).

In the following, when using the term ‘silicon carbide’ or ‘SiC’, we will always refer to the cubic (beta) modification of this polymorphous solid. Due to its symmetry, β -SiC has only one predicted infrared active mode, which is located at 796 cm^{-1} (position of the maximum of ϵ''), producing, in the case of small spherical particles, a maximum of its absorption efficiency factor at 935 cm^{-1} or $10.7\text{ }\mu\text{m}$. (For elongated ellipsoidal particles, this maximum of the absorption efficiency is shifted towards longer wavelengths: see Mutschke et al. 1999 and below.) As already explained, secondary emission bands of SiC can occur in the case of a non-ideal crystal structure, e.g. due to carbon or nitrogen inclusions. The case for N-doped SiC, for which Speck & Hofmeister refer to Suttrop et al. (1992), is problematic, however, since Suttrop et al. did not only detect a $20.9\text{ }\mu\text{m}$ (478 cm^{-1}) band, but also stronger additional bands in N-doped SiC. These additional bands (e.g. at 667 cm^{-1}

($15\text{ }\mu\text{m}$) and 580 cm^{-1} ($17.2\text{ }\mu\text{m}$)) are not seen in the sources of the astronomical ‘21’ μm feature. Moreover, the whole series of nitrogen-induced secondary features of SiC is weaker than the SiC main feature at least by a factor of 1000. Therefore, very low temperatures (below 80 K) would be necessary to reverse this ratio of band strengths. According to Hrivnak, Volk & Kwok (2000), the dust temperatures at the *inner* radii of the dust shells of the ‘21’ μm sources are between 165 and 220 K; due to the relatively small ratio between outer and inner shell radius, the *average* temperature of the dust grains producing the ‘21’ μm will hardly be below 80 K even for very transparent dust species (see below, Fig. 11).

As for C-doped SiC, there is presently too little information at hand about its potential secondary bands. As long as no studies on analytically well-characterized SiC samples with carbon impurities are available, it remains unclear whether or not the properties of N-doped SiC can be extrapolated to C-doped SiC.

Interestingly, there is yet another possibility to produce an 11 plus a ‘21’ μm band by, so to speak, ‘SiC plus something’: namely by coating SiC grains with SiO_x .

Recently performed experiments (Clément et al. 2003) demonstrate that pure SiC nanoparticles, under the influence of the laboratory atmosphere, are quite easily oxidized at their surfaces. This effect is shown in Fig. 2, where we compare the transmittance of SiO_2 grains with the transmittance of partially oxidized SiC nanoparticles (both embedded in KBr). The method for the production of SiC nanoparticles by laser pyrolysis and the in-




Fig. 2.— Comparison of the transmission spectrum of partially oxidized nano-SiC particles with that of amorphous SiO₂ grains.




Fig. 3.— Core-mantle particle consisting of SiC and amorphous silicon oxide. The silicon carbide core is identified by its characteristic lattice fringes.




Fig. 4.— Commercial β -SiC (dotted line) was heated in air at 1050° C (dashed) and has afterwards been treated with HF, thereby returning from an almost completely oxidized state (carrying a 9 μ m and a strong 21–22 μ m absorption band) to its original state characterized by the well-known 11 μ m band. The additional broad 7 μ m and 14 μ m absorptions in SiC are due to plasmon-phonon coupling, see Mutschke et al. (1999) for details.




Fig. 5.— Volume normalized absorption efficiency Q_{abs}/a of core-mantle particles composed of an SiC core and an amorphous SiO₂ mantle, the size of the whole (spherical) grains being small compared to the wavelength.

frared spectra of these grains have been discussed extensively by Clément et al. (2003). They also report the detection of an amorphous silicon oxide mantle by means of transmission electron microscopy (see Fig. 3). The large surface-to-volume ratio of the SiC core favours the formation and growth of a silicon oxide mantle, which can be a few nanometers thick and thus reach a considerable volume fraction of the particulate. Further evidence for the formation of SiO₂ on SiC surfaces, has been delivered – even though for significantly higher temperatures – by Mendybaev et al. (2002) and Tang & Bando (2003).

Depending on the particle size and shape as well as agglomeration effects, SiC nanoparticles show one strong IR resonance band, located at 10.8–12.4 μm. However, if partially oxidized at their surfaces, the SiC nanograins produce additional infrared bands at 9 and 21–22 μm. These bands can be assigned to stretching and bending vibrations of the SiO₄ tetrahedra. The minor 12.5 μm band of SiO₂ is not detected in the spectrum of partially oxidized SiC nanoparticles, since it almost coincides with the SiC main band shifted to longer wavelengths by shape effects.

An additional proof for the oxidation of SiC is delivered by treating the oxidized grains with HF, which quickly reduces the oxidized grains, leaving only the pure silicon carbide cores. The spectroscopic result of this process is depicted in Fig. 4.

Not only experimentally, but also by calculating the absorption efficiency of core-mantle-particles composed of SiC and amorphous SiO₂, absorption spectra with the principal structure of those shown in Figs. 2 and 4 can be derived. For this pur-

pose, the optical constants of amorphous SiO₂ compiled in the first volume of Palik (1985-98) have been used. We performed a Lorentz fit to these optical constants in order to analyse the underlying oscillator parameters. The damping constants of the SiC oscillator has been set to 50 cm⁻¹ and that of the 456 cm⁻¹ oscillator of SiO₂ to 42 (instead of 27), assuming that the nanocrystalline structure is comparable to that of defect-bearing polycrystalline films (see Mutschke et al. (1999), Sect. 2.3). As for the 456 cm⁻¹ SiO₂ oscillator, a larger damping constant than that corresponding to the Palik data is justifiable also on the basis of the spectra published by Lehmann, Schumann & Hübner (1984). These authors studied the transmittance of thin SiO_x films (with x<2) and found a strong broadening of the 20–21 μm band with decreasing x. This result is of potential relevance for the coating of SiC by silicon oxide in carbon-rich circumstellar shells, since in these environments, the formation of oxygen-deficient SiO₂ is very well conceivable. For a summary of all Lorentz parameters used, see Tab. 1.

Fig. 5 shows the absorption efficiency factor Q_{abs}/a of spherical core-mantle-particles composed of SiC and amorphous SiO₂ with core volume fractions ranging from 40% to 60%. Three prominent maxima of Q_{abs} are discernible: a peak at 8.3 μm, which originates from the SiO₂ mantle; the sharp 11 μm SiC resonance; and, finally, a 20 μm peak, with a main peak and a shoulder at the long-wavelength side, originating again from the SiO₂ mantle. Peak splitting is typical for bands of mantle materials. It arises from the existence of two interfaces with core and am-

Table 1: Frequencies (TO_j), oscillator strengths (Ω_j) and damping constants (γ_j) of SiC and SiO₂. We adopted the values $\epsilon_\infty = 6.49$ for SiC and $\epsilon_\infty = 2.2$ for amorphous SiO₂.

	TO_j (cm ⁻¹)	Ω_j (cm ⁻¹)	γ_j (cm ⁻¹)
SiC	796	1423	50
SiO ₂	456	417	42
	798	150	30
	1080	866	60
	1194	257	100
	1479	200	80

bient medium. All the bands are sharper than those of the transmission spectrum of partially oxidized SiC shown in Fig. 2. This difference in the bandwidth is primarily due to shape effects.

Looking at Fig. 5, one may assume that the depicted pattern of C_{abs}/V maxima is incompatible with the spectra of astronomical objects like HD 56126, since the 20 μm band, in this figure, is only a minor peak compared to those at 8.2 and 11 μm. However, such a conclusion would be premature given that the dust in the sources of the ‘21’ μm feature is rather cold, as has been highlighted in Sect. 2. For dust particles with temperatures below 150 K, all features at wavelengths smaller than ~20 μm are strongly attenuated, because they tend to be located in the Wien regime of the corresponding Planck functions. This effect, which was already noted by Speck & Hofmeister (2004), is illustrated in Fig. 6.

It is evident from this figure that cold core-mantle-particles composed of SiC and




Fig. 6.— Products of the absorption efficiency for SiC-SiO₂ core-mantle particles (core volume fraction = 0.5) with two different Planck functions, corresponding to dust temperatures of 160 K and 100 K. In grey colour, the case of ellipsoidal particles (shape distribution according to Ossenkopf, Henning & Mutschke 1992) is overplotted.




Fig. 7.— Normalized emission spectrum of a dust shell composed of SiC-SiO₂ core-mantle grains. The shell is supposed to follow a $1/r^2$ density distribution and a temperature distribution resulting from the condition of radiative equilibrium (see appendix) and a certain geometry of the shell of HD 56126 as inferred from observational data (see below, sect. 3.3). The residual emission of HD 56126 is shown for comparison.

amorphous SiO_2 do produce a strong $20\mu\text{m}$ feature. There is a rather good agreement between the observed and the calculated band positions (see also Tab. 3 for a quantitative assessment), and the weakness of the observed $11\mu\text{m}$ feature can be explained by dust temperatures significantly lower than 160 K on the average. There remains a problem, however, with the *width* and *shape* of the $20\mu\text{m}$ band produced by SiO_2 . For spherical particles, the FWHM of the $20\mu\text{m}$ SiO_2 emissivity maximum amounts to $1.9\mu\text{m}$, compared to an observed FWHM of $2.2\text{--}2.3\mu\text{m}$. Furthermore, the shoulder at the ‘red’ wing of the $20\mu\text{m}$ SiO_2 feature has not been seen in the astronomical ‘ $21\mu\text{m}$ ’ band. The bandwidth discrepancy can be overcome by a CDE; the shoulder at the ‘red’ wing of the band, however, does not vanish in the CDE case (see Fig. 7); nor does it vanish by any particular (realistic) choice of the dust temperature or dust temperature distribution.

3.2. Oxidized silicon grains

Instead of growing on top of silicon carbide grains, SiO_2 may equally well form mantles on top of pure silicon particles. Li & Draine (2002) were the first to examine the spectra emerging from silicon grains with oxide coatings, focussing on the case of stochastically heated nanoparticles. Smith & Witt (2002) proposed silicon nanoparticles as carriers of the so-called *extended red emission* observed in circumstellar environments; however, this identification is still a matter of debate.

The principal structure of the IR spectrum of Si-SiO₂ core-mantle-grains is similar to that of SiC-SiO₂ core-mantle-grains,

except for the absence of the $11\mu\text{m}$ band. In addition to the $8.2\mu\text{m}$ and the $20\mu\text{m}$ peaks, a rather strong $12.5\mu\text{m}$ band appears in the Si-SiO₂ absorption efficiency spectrum. The exact position λ_{20} of the $20\mu\text{m}$ resonance depends on the volume fraction f of the Si core. λ_{20} increases from 19.7 to $19.8\mu\text{m}$ as f decreases from 0.6 to 0.4 .




Fig. 8.— Products of the volume normalized absorption efficiency C_{abs}/V shown in the previous figure with two different Planck functions, corresponding to dust temperatures of 160 K and 100 K .

Following the procedure that has been explained in the previous subsection, we folded the absorption efficiency of a Si-SiO₂ core-mantle-grain with $f=0.5$ with two different Planck curves $B_\nu(T_d)$. For the dust temperature, the two values $T_d=100\text{ K}$ and $T_d=160\text{ K}$ have been chosen. As for the $20\mu\text{m}$ band, the result –

plotted in Fig. 8 – strongly resembles the case of SiC cores, which is not surprising, since the optical constants of the core have only a limited influence on the structure of those bands which are due primarily to the mantle.

It must be stressed, however, that core-mantle particles composed of Si and SiO₂ are more unlikely to be the carrier of the ‘21’ μm band than those composed of SiC and SiO₂, since it is questionable whether *silicon* grains can be formed in carbon-rich PPNe at all.

3.3. Cold FeO (wustite)

As mentioned in §1, iron oxides have been considered as carriers of bands in the 20–21 μm region by various authors already. It seems clear, though, that none of the ‘higher’ iron oxides like Fe₂O₃ or Fe₃O₄ can survive in a reducing environment. Iron monoxide (FeO, also called wustite²), by contrast, may survive in a carbon-rich circumstellar shell.

According to Duley (1980), the oxidation of metallic iron is a very efficient process provided that enough O₂ molecules are present. The degree to which iron particles are oxidized depends on the physical conditions, inter alia on the temperature. The *higher iron oxides* Fe₃O₄ or Fe₂O₃ become stable at higher temperatures (see, e.g., Gail & Sedlmayr 1998). By contrast, as Roberts (1961) and Fehlnert & Mott (1970) pointed out, several monolayers of *iron monoxide* form very quickly on top of Fe surfaces, even at low temperatures,

since this process requires a negligible activation energy. In order to end up with dust grains composed essentially of FeO (instead of a large metallic iron core and a tiny FeO mantle), the (low-temperature-)oxidation of *very small* iron grains is required – i.e. grains composed of 10³ atoms at maximum. Only such small iron grains – their diameter amounts to ~ 2 nm – could be fully converted into FeO by the process mentioned above. The oxidation of iron grains consisting of less than 10³ atoms is to some extent an artificial assumption; less artificial, though, than the constraint that the TiC clusters – which have previously been considered as carriers of the ‘21’ μm band – should consist of no more than 125 atoms to efficiently produce the TiC cluster signature.

Why would FeO, according to our suggestion, not exist in the vast majority of the PNe, at least not with high number densities? A plausible explanation is that in PNe with their large fractional ionization and high UV photon densities, iron oxides will be reduced to metallic iron. While the non-detection of the ‘21’ μm band in PNe spectra can thus be accounted for, its non-detection in the spectra of most of the oxygen-rich as well as carbon-rich AGB stars is more difficult to understand within the iron grain oxidation scenario for PPNe. If in PPNe like HD 56126, FeO forms by oxidation of iron, why does the same process not take place in C-stars, S-stars and in the vast majority of M-AGB stars? This may be due to the fact that the dust around these stars is, on the average, much hotter than dust in PPNe and that the sticking probability of oxygen on iron decreases as the temper-

²Strictly speaking, wustite is the terrestrial form of iron monoxide and has the chemical composition Fe_{1-x}O with $x = 0.05 \dots 0.16$.

ature increases. The Mg-Fe-oxide formation around some M-stars, postulated by Posch et al. (2002), must involve an entirely different mechanism than the low-temperature oxidation of iron in PPNe. Indeed, Ferrarotti & Gail (2003) predict Mg-Fe-oxides via the condensation of the very refractory MgO in M-stars. It is noteworthy that HD 56126 and other sources of the ‘21’ μm band studied by van Winkel & Reyniers (2000) are depleted in metals including iron. Other PPNe do not show the feature, even though more iron is available for FeO formation. There could be two reasons for that. One is that the iron particles should be small as stated above. The second is that FeO formation is essentially governed by oxygen availability and that the iron abundance is therefore only crucial if there is enough free oxygen.

Whereas the knowledge on the Fe-Mg-oxide formation in circumstellar environments is still scarce, quite detailed *spectroscopic* information on Mg-Fe-oxides is available. The optical constants of $\text{Fe}_x\text{Mg}_{1-x}\text{O}$ samples (x amounting to 0.4 ... 1.0) with magnesiowustite structure have been measured at room temperature by Henning et al. (1995). Subsequently, Henning & Mutschke (1997) published *low temperature data* for various cosmic dust analogs, i.e. their indices of refraction $n(\lambda)$ and absorption $k(\lambda)$ for $T = 10, 100, 200$ and 300 K. A major change of the infrared spectrum of $\text{Fe}_x\text{Mg}_{1-x}\text{O}$ is observed as the samples are cooled: namely a sharpening of the vibration band located at $18\text{--}20\text{ }\mu\text{m}$ (depending on x). Fig. 9 shows this effect for $x=1.0$, i.e. for pure FeO. (The case $T=10$ K has been omitted, since it does not significantly differ from $T=100$ K.)




Fig. 9.— Absorption efficiency factors C_{abs} of spherical FeO grains, calculated from their optical constants measured at room temperature ($T = 300$ K) and at low temperatures ($T = 200$ K, $T = 100$ K). The respective C_{abs} functions have been multiplied with Planck functions $B_\nu(T)$ (normalized at $20\text{ }\mu\text{m}$) for the appropriate temperatures.

The FWHM of the $20\mu\text{m}$ FeO band decreases from $3.6\mu\text{m}$ at room temperature to $2.4\mu\text{m}$ at $T = 100\text{ K}$, while the band position is shifted from $\lambda_{\text{max}} = 19.9\mu\text{m}$ to $\lambda_{\text{max}} = 20.1\mu\text{m}$. The latter effect is exclusively due to the multiplication with the Planck functions $B_\nu(T)$, which have maxima positions of 17, 25.5 and $51\mu\text{m}$ for $T = 300, 200$ and 100 K , respectively. The rather large width of the $20\mu\text{m}$ band produced by FeO, in combination with the fact that for $T = 200\text{ K}$ and $T = 100\text{ K}$, the peak position is located at the ‘blue’ wing of the corresponding Planck curve, leads to an asymmetric profile of the $C_{\text{abs}} \times B_\nu(T)$ curves – the same kind of asymmetry which is found for the astronomical ‘ $21\mu\text{m}$ ’ band.

Iron monoxide has a rather high absorption efficiency in the visual range of the spectrum. Therefore, it is comparatively strongly heated in the radiation field of an F or G star. An equilibrium temperature of less than 100 K is reached only at distances from the central star larger than 2×10^4 stellar radii. This can be shown by calculating the energy balance for the case of an equilibrium between the radiation field of the central star and the dust grains in its circumstellar according to the scheme summarized in the appendix.

According to Kwok, Volk & Hrivnak (2002), the inner radius R_{in} of the dust shell of IRAS 07134+1005 is located at $1.2''$ from the central star, corresponding to $1.2 (1.8) \times 10^{-2}\text{ pc}$ for an assumed distance of 2 (3) kpc (Kwok, Volk & Hrivnak (2002) assume a distance of 2 kpc, Meixner et al. (2003) assume 3 kpc). In terms of the stellar radius corresponding to a luminosity of $1400 L_\odot\text{ kpc}^{-2}$ (Hrivnak, Volk & Kwok 2000) and a temperature of 7000 K ,

this amounts to 9760 or 6490 stellar radii for $d=2$ (3) kpc, respectively.




Fig. 10.— Radiative equilibrium temperature of FeO in the radiation field of a star with an effective temperature of 7000 K (solid line). For comparison, the radial temperature profile dust with wavelength-independent optical constants is also shown (dashed line; see also the appendix to this paper). Short-dotted lines delimit the r -range of the brightest part of IRAS 07134+1005.

In Fig. 10, the position of $R_{\text{in}}/R_{\text{eff}} = 6490$ has been highlighted by a dotted vertical line. The equilibrium temperature of FeO corresponding to this distance from the central star amounts to 155 K . This is in good agreement with the results of the model calculations by Hrivnak, Volk & Kwok (2000), who derive an inner temperature of the dust shell of 165 K by radiative transfer calculations aimed at a fit of the

spectral energy distribution.




Fig. 11.— Equilibrium temperatures of amorphous carbon, FeO and core-mantle particles composed of SiC and SiO₂ in the radiation field of an F5 star. The domain from which presumably most of the IR emission of IRAS 07134+1005 comes is indicated by dotted lines.

In terms of the ratio between outer and inner shell radius, the dust shells of the PPNe showing the ‘21’ μm feature are rather thin. For IRAS 07134+1005, the observations of Kwok, Volk & Hrivnak (2002) and Miyata et al. (2003) yield an outer shell radius not much larger than twice the inner shell radius. This implies that the dust temperature range, for a $r^{-0.4}$ -like decrease of $T(r)$,³ is rather small. For FeO, we derive a dust tem-

perature at $2 R_{\text{in}}$ of 120 K, meaning that $120 \text{ K} < T_d < 155 \text{ K}$ (again for $d=3 \text{ kpc}$). For other dust species which are generally assumed to be present in carbon-rich PPNe, narrow T_d ranges are found as well. Amorphous carbon is warmer than FeO due to the monotonous increase of its absorption efficiency factor from the IR to the UV: we find $153 \text{ K} < T_d < 200 \text{ K}$ under the same conditions as above; core-mantle particles composed of SiC and SiO₂, unless heavily N-doped, remain cooler than FeO ($104 \text{ K} < T_d < 126 \text{ K}$;⁴ see Fig. 11).

These estimates make it possible to calculate the emission spectrum of an optically thin dust shell with a homogeneous density distribution and a temperature range of $120 \text{ K} < T_d < 155 \text{ K}$. If we are only interested in the *profile* of the emerging spectral energy distribution, it suffices to calculate a mean of blackbody functions $B_\nu(T)$ for $120 \text{ K} < T_d < 155 \text{ K}$ and to multiply the result with $C_{\text{abs}}(T_m)$. The resulting normalized spectrum is shown in Fig. 12. Both with respect to feature position and width, the emission spectrum of FeO calculated for the temperature range indicated above reproduces the observed ‘21’ μm band profile fairly accurately, even though no *ad hoc* assumptions concerning the dust temperature have been made.

For a CDE, quite different spectra of FeO emerge. For a CDE according to Ossenkopf, Henning & Mathis (1992) (‘CDE-Osk’; it is characterized by a maximum probability for particle shapes close to the spherical shape), the position of the FeO absorption efficiency maximum is shifted

³See the appendix for a comment on the meaning of the $T(r) \propto r^{-0.4}$ decrease.

⁴These values refer to a core volume fraction of 0.5; the larger the (rather transparent) SiO₂ mantle, the lower the dust temperature will be.




Fig. 12.— Normalized emission spectrum of an FeO dust shell (solid: spherical particles, dotted: CDE according to Ossenkopf et al. (1992)) with a temperature distribution corresponding to the range highlighted in Fig. 11 and a $1/r^2$ density distribution. For comparison, the residual emission of IRAS 07134+1005 is shown.

to $20.7\text{--}21\text{ }\mu\text{m}$, and the bandwidth changes even more dramatically, i.e. it increases to $6.5\text{--}7.6\text{ }\mu\text{m}$ (see Fig. 12). For a CDE with equal probability of all particle shapes, the changes in feature position and shape are even more dramatic, but this kind of shape distribution is unlikely to represent the shapes of circumstellar grains. The assumption that nearly spherically symmetric grains dominate in circumstellar dust shells is more reasonable, hence the ‘truth’ should lie between the ‘CDE-Osk’ case and the case of spherical particles.

Fig. 12 may give the impression that FeO dust can produce a feature comparable to the astronomical ‘ $21\text{ }\mu\text{m}$ ’ emission band only under *very* particular physical conditions, e.g. only for $T < 155\text{ K}$, while we know from Hrivnak, Volk & Kwok (2000) that the dust temperatures in the sources of the ‘ $21\text{ }\mu\text{m}$ ’ band can reach 220 K . However, as long as $T < 250\text{ K}$, $B_v(T)$ will always have its maximum at a wavelength larger than $20\text{ }\mu\text{m}$, thus shifting the $19.9\text{ }\mu\text{m}$ emissivity maximum of FeO to a feature position of $20.0\text{--}20.1\text{ }\mu\text{m}$. And indeed, $T < 250\text{ K}$ seems to be fulfilled for all known sources of the ‘ $21\text{ }\mu\text{m}$ ’ feature. In a wider sense, however, it is true that FeO can produce a strong $20.1\text{ }\mu\text{m}$ feature only under particular physical conditions. If the mean dust temperature is much smaller than 100 K , too little energy will be contained in the $20\text{ }\mu\text{m}$ FeO band to produce a feature as strong as that observed in some PPNe. If, by contrast, $T > 250\text{ K}$, the feature will become too broad and too much shifted to the ‘blue’ as to account for the astronomical ‘ $21\text{ }\mu\text{m}$ ’ band.⁵

⁵Henning & Mutschke (1997) point out that only

3.4. Remark on the elemental abundances required to account for the ‘21’ μm band

Li (2003) made a very useful estimate of the mass of dust which is required to account for the radiative energy contained in the ‘21’ μm band in the case of IRAS 07134+1005. He pointed out that the Kramers-Kronig equation relates the integral over the absorption efficiency of a grain to its volume and the orientational averaged polarizability F . Since for all materials with a static dielectric constant >4 and for moderately elongated grain shapes, the latter quantity has a value between 0.5 and 1 (Purcell 1969, Fig. 1), there is an upper limit to the integrated absorption efficiency per unit volume, which is more or less the same for the dust species considered here. Consequently, there is also a general upper limit to the energy emitted in the 21 μm band per unit particle mass or, vice versa, a minimum total grain volume for a certain total energy emitted through this band. Hence, the minimum mass derived by Li can be adopted to other materials just by scaling with the mass density of the grain material. This leads to the following conclusion concerning the abundance problem: While there is, according to Li (2003), about 100 times too little amount of titanium to account for the ‘21’ μm band by assigning it to TiC , there is 2.5 times more Fe than would be needed to account for the ‘21’ micron band with FeO (Fe is 250 times more abundant than Ti in HD 56126). Sulfur is 312 times more

abundant than Ti in the same star, silicon 1225 times, carbon 34500 times and oxygen 36000 times. Hence, compounds of iron, sulfur, silicon, carbon and oxygen cannot be excluded as carriers of the ‘21’ μm band in HD 56126 and objects with comparable elemental abundances.

4. Discarded feature carriers

For the sake of completeness, two potential carriers of the ‘21’ μm band which have been considered as promising candidates (but should no longer be considered) will be discussed: silicon disulfide and titanium carbide. In the case of TiC , insurmountable problems with the abundance of titanium exist, whereas SiS_2 can be ruled out by spectroscopic arguments.

4.1. Silicon disulfide

As mentioned in §1, the formation of sulfides is very likely in carbon-rich circumstellar environments. Indeed, the 26 μm feature, observed in many sources of the ‘21’ μm feature, is currently ascribed to magnesium sulfide (MgS , Hony et al. 2003).

However, from a spectroscopic point of view, there are problems with ascribing the 20.1 μm band to SiS_2 . The main point is that this material has *two* infrared bands, of which the second, located at 16.8 μm for spherical particles (see Fig. 13), has no counterpart in the astronomical spectra. Furthermore, the position and FWHM of the main SiS_2 band, which amount to 19.8 and 1.6 μm , respectively, do not agree with the observed characteristics of the 20.1 μm band. For a CDE, this discrepancy becomes smaller

a limited number of cosmic dust analogs have optical constants which change as strongly at low temperatures as those of FeO (and FeS).

($\lambda_{\max} = 20.2 \mu\text{m}$, $\text{FWHM} = 1.8 \mu\text{m}$), but the problem with the unobserved secondary feature at $16.8 \mu\text{m}$ becomes even worse, given that the ratio between ~ 17 and $\sim 20 \mu\text{m}$ band increases for a CDE. Very low dust temperatures (significantly below 100 K) would be required to make the $\sim 17 \mu\text{m}$ band negligible in strength compared to the $\sim 20 \mu\text{m}$ SiS_2 band.

The visual and near-infrared optical constants of SiS_2 have not yet been derived, such that it is not possible to calculate $T(r)$ for SiS_2 based on its radiative energy balance. However, it is hardly conceivable that SiS_2 is so transparent in the visual range as to remain much colder than 100 K over the whole range indicated in Fig. 11.

4.2. Titanium carbide

Even though recent papers by Li (2003), Chigai et al. (2003) exclude TiC nanoparticles as the carrier of the ‘ $21 \mu\text{m}$ ’ band, we append a discussion of TiC ’s infrared properties here, focussing on the relation between bulk TiC and TiC clusters.

Macroscopic TiC has metallic properties, with an absorption index k which is larger than the refractive index n throughout the whole infrared range. For $\lambda > 13 \mu\text{m}$, both n and k are larger than 10. At about $19 \mu\text{m}$, a resonance peak occurs, which is rather weak, however, with a relative increase of n , k and C_{abs} by less than 10%.

Small TiC clusters, e.g. those composed of $4 \times 4 \times 4$ atoms, behave differently from bulk matter insofar as the mobility of the cluster electrons is reduced. The lattice resonance is then no longer a small additional term in the dielectric function, but the dominant contribution at infrared




Fig. 13.— Normalized emission spectrum of spherical SiS_2 particles for two different dust temperatures, 100 and 160 K. The $16.8 \mu\text{m}$ secondary emission band, which is not observed in the sources of the ‘ $21 \mu\text{m}$ ’ band.

frequencies. Therefore, a comparatively strong $20\text{ }\mu\text{m}$ emission band arises from the clusters; its exact position, however, depends on the cluster size. Chigai et al. (2003) derived the following Lorentz oscillator parameters for the small TiC clusters produced by von Helden et al. (2000): a resonance frequency of 488 cm^{-1} , a damping constant of 50.2 cm^{-1} and an oscillator strength of 187 cm^{-1} . For spherical particles, this translates into a maximum position of the absorption efficiency factor of $20.0\text{ }\mu\text{m}$ and an FWHM of $2.0\text{ }\mu\text{m}$. For a CDE, no shift of the band position is found in this case (due to the non-negative values of the dielectric function of TiC according to Chigai et al. (2003)), but only an increase of the FWHM to $2.2\text{ }\mu\text{m}$ (for a CDE according to Ossenkopf, Henning & Mathis 1992).

Fig. 14 shows the normalized emission of a small TiC cluster (at $T_d=160\text{ K}$), the normalized dust emission arising from macroscopic dust grains (at two different temperatures $T_d=160\text{ K}$ and $T_d=120\text{ K}$) and, again for comparison, the profile of the astronomical ‘21’ μm feature as derived from the ISO-spectrum of IRAS 07134+1005. Since the maximum of the absorption efficiency factor of TiC is not evident from Fig. 14 due to the normalization, it should be added that C_{abs}/V (TiC) does not exceed $0.6\text{ }\mu\text{m}^{-1}$. (Depending on the value of ϵ_∞ , C_{abs}/V may also be substantially smaller). By contrast, the $\sim 20\text{ }\mu\text{m}$ peak absorption efficiency of cold (100 K) FeO as well as that of amorphous SiO_2 amounts to $1.6\text{ }\mu\text{m}^{-1}$.

5. Conclusions

Tables 2 and 3 summarize the spectro-




Fig. 14.— Normalized emission spectrum of bulk TiC at temperatures of 120 K (dashed) and 160 K (solid) and of TiC nanoparticles at $T = 160\text{ K}$ (dotted) compared to the residual emission of IRAS 07134+1005.

scopic information on the potential carriers of the ‘21’ μm feature in two different respects. Tab. 2 gives an overview of the positions of the absorption efficiency maxima and of the corresponding bandwidths. Tab. 3, on the other hand, refers to normalized dust emission spectra like those shown in Figs. 7, 12, 13 and 14, hence to products of the absorption efficiency with blackbody functions. These products are compared with the repeatedly shown profile of the ‘21’ μm band: we present the sums (henceforth: $\Sigma\delta^2$) of the squares of the differences between the ‘profile’ and the ‘dust emission’ values, for wavelengths between 17 and 25 μm (outside this range, the ISO profile is characterized by comparatively large noise). The ‘temperature grid’ has been chosen such as to cover the range between 160 K and 70 K in steps of 10 K. The bold numbers indicate the respective minima of $\Sigma\delta^2$. For SiC+SiO₂-sph, TiC-sph and FeO-sph, the minimum of $\Sigma\delta^2$ occurs at blackbody temperatures between 80 and 90 K. This is due to the asymmetric shape of the profile of the ‘21’ μm band. On the basis of a symmetric absorption efficiency profile, this asymmetric profile is best reproduced by multiplication with a blackbody of 80–90 K. If, however, the absorption efficiency of the feature carrier has an *intrinsically* asymmetric profile (as for ellipsoidal SiC+SiO₂-core-mantle grains), it is not necessary to assume such a low dust temperature (120 K are sufficient in this case). The $\Sigma\delta^2$ values for SiS₂ are particular insofar as they continuously decrease with decreasing temperature, down to less than 70 K. This has a very simple reason: The 16.8 μm secondary band of SiS₂ (see Fig. 13), which has no counterpart in the astronomical spectra, is the more sup-

pressed the lower the temperature. As regards the absolute values of $\Sigma\delta^2$, they are smallest for TiC clusters and second smallest for ellipsoidal SiC+SiO₂ core-mantle grains.

Despite the good spectroscopic fits obtainable with TiC, cold SiC+SiO₂ and cold FeO, we come to the conclusion, based on the properties of the sources of the ‘21’ μm band, that none of the hitherto discussed candidate carriers is a convincing feature carrier candidate in all relevant respects. Even though there are oxides which may appear as likely band carriers from a spectroscopical point of view, it is difficult to conceive of a scenario accounting for their formation in carbon-rich PPNe. On the other hand, dust species which are usually predicted to form in carbon-rich environments, like carbon, carbides and sulfides do either not match the observed ‘21’ μm band profile (e. g. SiS₂), or even their maximum abundance is too small for emitting the energy contained – at least in some cases – in the ‘21’ μm feature (which rules out TiC). As for the individual groups of feature carriers, our arguments are the following ones:

- 1) Dust and macromolecular species which produce a 20–22 μm emission by transitions involving heteroatoms in carbonaceous networks (e.g. Papoular 2000), are not likely to be the feature carrier due to the uniqueness of the ‘21’ μm band profile on the one hand and the non-uniqueness of the heteroatom-caused emission bands on the other hand.

- 2) There do exist oxide dust species which can produce a 20.1 μm emission band very effectively (if the question of their formation in C-rich environments is

Table 2: Overview of the most promising carriers of the ‘21’ μm feature which have been discussed in the present paper. Note: The observed position and FWHM of the ‘21’ μm band are 20.1 μm and 2.3 μm , respectively. ‘sph’ means spherical particle shape, ‘CDE-Osk’ designates a continuous distribution of ellipsoids according to Ossenkopf, Henning & Mathis (1992). The λ ($C_{\text{abs,max}}$) values do *not* directly represent feature positions, since these result from the product $C_{\text{abs}} \times B_{\nu}(T_{\text{d}})$; for dust temperatures close to 150 K, this leads to a shift towards the ‘red’ of 0.1–0.2 μm .

candidate carrier	particle shape	λ ($C_{\text{abs,max}}$) [μm]	FWHM (20) [μm]
SiC+SiO ₂	sph. ^(a)	8.1–8.3, 11.0–11.1, 12.5, 19.7–19.8	1.9
	CDE-Osk ^(a)	8.2–8.4, 11.1–11.2, 19.8–20.0	2.1–2.6
Si+SiO ₂ ^(a)	sph.	8.0, 12.5, 19.7–19.8	1.8
FeO ^(b)	sph.	19.9	2.3
	CDE-Osk	20.7	6.5
SiS ₂	sph.	16.8, 19.8	1.6
	CDE-Osk	16.8, 20.2	1.8
TiC clusters	sph	20.0	2.0
	CDE-Osk	20.0	2.2
TiC grains	sph	19.2	– ^(c)

^(a)core-mantle-particles with SiC core volume fractions between 0.4 and 0.6

^(b)at an average temperature of ~ 140 K

^(c)broad bump, FWHM not defined

Table 3: Selected potential carriers of the ‘21’ μm feature: quality of fits to the profile of the ‘21’ μm band for different *mean* dust temperatures T_d (between 130 K and 70 K). As an indicator for the quality of an individual fit, the sum of the squares of the differences ‘normalized 21 μm flux’ minus ‘ $C_{\text{abs}} \times B_\nu(T_d)$ ’ has been calculated for $17 \mu\text{m} < \lambda < 25 \mu\text{m}$. This sum corresponds to the usual ‘ χ^2 ’ insofar as the quantities subtracted from each other are already normalized.

T_d [K]	SiC+SiO ₂ -sph ^(a)	SiC+SiO ₂ -CDE ^(a)	FeO-sph	TiC-sph	SiS ₂ -sph
160	5.98	3.10	6.40	3.54	18.3
150	5.66	2.92	6.03	3.29	17.8
140	5.31	2.80	5.62	3.02	17.4
130	4.94	2.71	5.20	2.72	16.9
120	4.55	2.67	4.78	2.41	16.4
110	4.14	2.74	4.37	2.08	15.8
100	3.76	3.00	4.03	1.76	15.1
90	3.47	3.63	3.87	1.50	14.3
80	3.42	4.99	4.13	1.40	13.4
70	4.02	7.94	5.41	1.74	12.4

^(a)core-mantle-particles with a SiC core volume fraction of 0.5

left aside). As we have shown, this holds true both for cold FeO and for SiO₂. Iron monoxide has *only* one IR active mode, centered at 19.9 μm , which shows an exceptional narrowing at temperatures in the 200–100 K range. By contrast, amorphous SiO₂ has two strong IR active modes, one of them being centered at 20.5 μm . Our laboratory experiments show that amorphous SiO₂ easily forms mantles on top of SiC. Core-mantle particles composed of SiC and SiO₂, if colder than 150 K, emit mainly through the lower energy band of SiO₂. Thus they can in principle account for the ‘21’ μm band, even though the reproduction of the feature profile is not as good as in the case of cold FeO.

But there is a major problem with any scenario involving oxides as band carriers:

the difficulty to explain how oxides can form or at least survive in carbon-rich PPNe. It is hardly conceivable that the carrier of the ‘21’ μm band is just a remnant of the oxygen-rich outflow phase of the respective former AGB stars. The interval between the oxygen-rich phase and the carbon-rich PPN phase is too long as to account for a co-location, as proven for HD 56126, of products of the C-rich and the O-rich phase. The remnants of the O-rich phase are expected to be located at least one order of magnitude further out on a radial scale than the products of the C-rich phase. How, then, can either SiC be oxidized or FeO form/survive in C-rich environments? As for the first process, there are several studies showing that *complete* transformation of SiC to SiO₂ occurs

at temperatures above 700 K (e.g. Mendybaev et al. 2002, Tang & Bando 2003); provided that large defective surfaces as those of nanograins are available for the reaction, the formation of SiO₂ mantles on top of SiC grains also takes place at room temperature as our laboratory experiments indicate (see Fig. 3). But why should this SiO₂ mantle formation take place preferentially in PPNe with C/O ratios close to or larger than unity? This question, unfortunately, has to remain open here. As for the hypothesis that FeO is the carrier of the ‘21’ μ m band, it would require the oxidation of Fe to FeO in cold circumstellar environments. It seems that metal-poor PPNe like HD 56126 – paradoxically, at the first glance – provide good conditions for FeO formation, since this process requires the presence of small metallic iron grains. The *absence* of any feature attributable to FeO in the vast majority of the Planetary Nebulae, on the other hand, could be accounted for by the fact that FeO can persist only in a narrow range of equilibrium between oxidation (to higher iron oxides) and reduction (to metallic iron). Thus, in PNe, FeO is probably reduced to metallic iron.

3) Quantum size effects critically influence the optical properties of very small grains and clusters. Phonon modes can shift when the lattice changes from ‘infinite’ to finite size, and conductivity may be reduced. For the TiC *grainlets* proposed as carriers, this means that a ‘21’ μ m band would only occur for clusters and particles in a *very narrow size range*. This makes it unlikely that there would be a sufficient number of TiC particles in the PPNe to account for the observed band strength,

given that the predicted Ti abundance is not large enough and that mantling by amorphous carbon will reduce the band strength even further.

4) Any identification or ruling-out of a candidate carrier is strengthened if it does not only have one, *but two or more observable infrared bands*. If these bands are separated by $\geq 10 \mu$ m, the secondary bands may be suppressed by temperature effects. This is likely to occur for the 11 μ m SiC band expected to arise from the core-mantle particles mentioned above. However, in the case of SiS₂, even temperatures in the order of 100 K will not be sufficient to suppress the 16.8 μ m secondary emission feature. The multiband identification problem also sets constraints on bands produced by impurities, because the impurity-induced bands are usually much weaker than the phonon bands of the respective matrix materials (which provides a further argument against doped SiC as the carrier of the ‘21’ μ m feature).

We thank Gabriele Born, Jena, for preparing our samples. Ingrid Hodouš, Vienna, kindly provided us with the ISO-spectra of IRAS 07134+1005 and SAO 34504, reduced with the ISO spectral analysis package ISAP. An anonymous referee contributed some substantial arguments to the original version of this paper and thereby improved it. TP received a DOC grant from the Austrian Academy of Sciences. HM acknowledges support by DFG grant Mu 1164/5.

A. Appendix: On the temperature profile and spectral energy distribution of geometrically thin PPNe dust shells

In the previous sections, calculations of the radiative equilibrium temperature of the sources of the ‘21’ μm feature have been presented. The temperature profiles in turn have been partly used to calculate model spectra. We append a discussion of the method used for the dust temperature (T_d) calculation as well as of the influence of T_d on the emerging spectral energy distributions (SEDs).

In the radiative equilibrium case and for optically thin shells (where backwarming effects can be neglected), the radial variation of T_d is derived from the following energy balance:

$$\int_0^\infty \pi a^2 Q_{\text{abs}}(\lambda) \pi B_\lambda(\lambda, T_{\text{eff}}) \frac{R_{\text{eff}}^2}{R^2(T_d)} d\lambda = \int_0^\infty 4\pi a^2 Q_{\text{abs}}(\lambda) \pi B_\lambda(\lambda, T_d) d\lambda \quad (\text{A1})$$

The first term describes the energy absorbed by an individual dust grain with a radius a and an absorption efficiency factor $Q_{\text{abs}}(\lambda)$ in a stellar radiation field $B_\lambda(\lambda, T_{\text{eff}})$ (T_{eff} denoting the effective stellar temperature); the second term describes the energy emitted by the same dust grain. R_{eff} is the effective stellar radius, whereas $R(T_d)$ is the distance of the grain with temperature T_d from the star. While the above integral runs from zero to infinity, numerical integration can of course be performed only between a certain lower (λ_{min}) and a certain upper limit (λ_{max}). We chose $\lambda_{\text{min}} = 0.05 \mu\text{m}$ and $\lambda_{\text{max}} = 500 \mu\text{m}$. It is difficult, however, to derive the values of Q_{abs} close to λ_{min} ; some of the sets of optical constants have to be extrapolated to UV wavelengths since measurements of the UV properties of cosmic dust analogues are still lacking. For FeO, e.g., our measurements extend to $0.2 \mu\text{m}$; beyond this wavelength, n and k were assumed to be constant. Any other assumption, and even the choice $\lambda_{\text{min}} = 0.2 \mu\text{m}$, would not significantly change our results.

We recall that simple power-law models for $Q_{\text{abs}}(\lambda)$ make it possible to solve eq. (A1) analytically for $T(r)$. For $Q_{\text{abs}}(\lambda) \propto 1/\lambda$ – the case corresponding to small dust particles with wavelength-independent (practically: weakly wavelength-dependent) optical constants n and k –, $T(r)$ decreases with $1/r^{0.4}$. This solution to eq. (A1) has been plotted in Fig. 10. (For the less realistic case $Q_{\text{abs}}(\lambda) = \text{const.}$, $T(r)$ decreases with $1/r^{0.5}$.)

In the main part of our paper, it has been shown that the temperature range corresponding to the measured spatial extension of a typical PPN like HD 56126 is comparatively narrow for all the dust species examined. The ratio between the temperature at the outer and the temperature at the inner dust shell edge (i.e., its dense observable part) is typically in the order of 1.3. This has an important consequence for the calculation of the dust shell’s SED. Again in the optically thin case, the latter is given by

$$F_d(\lambda) = c \times Q_{\text{abs}}(\lambda) \int_{r=1}^{r=R_{\text{out}}/R_{\text{in}}} B_\lambda(\lambda, T_d(r)) dr, \quad (\text{A2})$$

where c is a product containing the dust column density, the average geometrical cross

section of the dust grains, and the surface of the inner dust shell boundary (e.g. Schutte & Tielens (1989)). If R_{out}/R_{in} is indeed in the order of 2 (as for HD 56126 according to the observations of Kwok, Volk & Hrivnak (2002) and Miyata et al. (2003)), the result of the integral on the right-hand side of eq. (A2) is surprisingly similar to a single black-body function for a temperature intermediate between $T_d(R_{in})$ and $T_d(R_{out})$. This effect is shown in Fig. 15, where we have set $c=1$ and $Q_{abs}(\lambda)=1$ for simplicity.




Fig. 15.— Spectral energy distributions ($F_\nu(\lambda)$) emerging from a single blackbody (dotted line, for $T=175$ K) compared to SEDs emerging from a sum of blackbody contributions as the one entering into eq. (A2). For $R_{out}/R_{in} \leq 2$ (solid and dash-dotted line), the blackbody-sum is virtually not distinguishable from a single blackbody with a temperature intermediate between $T(R_{in})$ and $T(R_{out})$. Only for large values of R_{out}/R_{in} (e.g. values close to 10000, as for the dashed line), the blackbody-sum has a profile distinctly different from that of a single Planck function.

It is evident from this figure that the the SED calculated for a geometrically thin circumstellar shell (using realistic absorption efficiencies) according to eq. (A2) depends much more on the order of magnitude of the dust temperature (i. e. on whether the mean dust temperature amounts to 175 or 125 K) than on actually calculating a weighted average of Planck functions.

On the other hand, even a reasonably good estimation of the mean dust temperature

$T_d(r)$ is difficult without any information on the values of $Q_{\text{abs}}(\lambda)$ in the visual and near-infrared range. For SiS_2 and TiC , e.g., no optical constants for wavelengths smaller than $10\,\mu\text{m}$ are available. Hence, the SEDs emerging from dust shells composed of these grain species can only be calculated by arbitrarily assuming mean T_d -values for them.

REFERENCES

- Buss, R.H., Cohen, M., Tielens, A.G.G.M., et al., 1990, *ApJ*365, L23
- Chigai, T., Yamamoto, T., Kaito, C., & Kimura, Y., 2003, *ApJ*587, 771
- Clément, D., Mutschke, H., Klein, R., & Henning, Th., 2003, *ApJ*594, 642
- Clément, D., Mutschke, H., Klein, R., Jäger, C., Dorschner, J., Sturm, E., & Henning, Th., 2004 *ApJ*, submitted
- Cox, P., 1990, *A&A*236, L29
- Dayal, A., Hoffmann, W.F., Biegging, J.H., Hora, J.L., Deutsch, L.K., & Fazio, G.G., 1998, *ApJ*492, 603
- Duley, W.W., 1980, *ApJ*240, 950
- Fehlner, F.P., & Mott N.F., 1970, *Oxid. Met.* 2, 59
- Ferrarotti, A.S., & Gail H.-P., 2003, *A&A*398, 1029
- Gail H.-P., Sedlmayr E., 1998, in: *Chemistry and Physics of Molecules and Grains in Space*, Faraday Discussions no. 109, The Faraday Division of the Royal Society of Chemistry, London, p. 303
- Goebel, J.H., 1993, *A&A*278, 226
- Grishko, V.I., Tereszchuk, K., Duley, W.W., & Bernath, P., 2001, *ApJ*558, L129
- Henning, Th., Begemann, B., Mutschke, H., & Dorschner, J., 1995 *A&AS*112, 143
- Henning, Th. & Mutschke, H., 1997, *A&A*327, 743
- Henning, Th., & Mutschke, H., 2001, *Spectrochim. Acta* 57, 815
- Hill, A.G.M., Jones, A. P., & d'Hendecourt, L.B., 1998, *A&A*336, L 41
- Hony, S., Waters, L.B.F.M., & Tielens, A.G.G.M., 2001, *A&A*378, L41
- Hony, S., Tielens, A.G.G.M., Waters, L.B.F.M., & de Koter, A. 2003, *A&A*402, 211
- Hrivnak, B.J., Volk, K., & Kwok, S., 2000, *ApJ*535, 275
- Justtanont, K., Barlow, M.J., Skinner, C.J., Roche, P.F., Aitken, D.K., & Smith, C.H., 1996, *A&A*309, 612
- Kimura, Y. & Kaito, C., 2003, *MNRAS*343, 385
- Kwok, S., Volk, K., & Hrivnak, B. J., 1989, *ApJ*345, L51
- Kwok, S., Volk, K., & Hrivnak, B. J., 2002, *ApJ*573, 720
- Lehmann, A., Schumann, L., & Hübner, K., 1984, *phys. stat. sol. B* 121, 505
- Li, A., & Draine, B., 2002, *ApJ*564, 803
- Li, A., 2003, *ApJ* 599, L45
- Mendybaev, R.A., Beckett, J.R., Grossman, L., Stolper, E., Cooper, R.F., & Bradley, J.P., 2002, *Geochim. Cosmochim. Acta*66, 661
- Meixner, M., Zalucha, A. Fong, D., & Justtanont, K., 2003, in: *Astrophysics of Dust*, ed. A.N. Witt, poster P73

- Miyata, T., Kataza, H., Okamoto, K.Y., et al., 2003, in: *Astrophysics of Dust*, ed. A.N. Witt, poster P74
- Mutschke, H., Andersen, A. C., Clément, D., Henning, Th., & Peiter, G. 1999, *A&A*345, 187
- Ossenkopf, V., Henning, Th., & Mathis, J.S., 1992, *A&A*261, 567
- Oudmaijer, R.D., & de Winter, D., 1995, *A&A*295, L43
- Palik, E. D. (ed.) 1985-1998, *Handbook of Optical Constants of Solids*, 3 vols. (Boston MA: Academic Press)
- Papoular, R., 2000, *A&A*362, L9
- Posch, Th., Kerschbaum, F., Mutschke, H., Fabian, D., Dorschner, J., & Hron, J., 1999, *A&A*352, 609
- Posch, Th., Kerschbaum, F., Mutschke, H., Dorschner, J., & Jäger, C. 2002, *A&A*393, L7
- Purcell, E. M. 1969, *ApJ*, 158, 433
- Roberts W.M., 1961, *Trans. Faraday Soc.* 57, 99
- Schutte, W.A., & Tielens, A.G.G.M., *ApJ*343, 369
- Smith, T.L., & Witt, A.N., 2002, *ApJ*565, 304
- Speck, A.K., & Hofmeister, A.N., 2004, *ApJ*600, 986
- Sourisseau, C., Coddens, G., & Papoular, R., 1992, *A&A*254, L1
- Suttrop, W., Pensi, G., & Choyke, W.J., Stein, R., & Leibenzeder, S., 1992, *J. Appl. Phys.* 72, 3708
- Tang, Ch., & Bando, Y., 2003, *Appl. Phys. Lett.* 83, 659
- Volk, K., Kwok, S., & Hrivnak, B.J., 1999, *ApJ*516, L99
- Volk, K., Xiong, G.-Z., & Kwok, S., 2000, *ApJ*530, 408
- Van Winckel, H., & Reyniers, M., 2000, *A&A*354, 135
- Von Helden, G., Tielens, A., van Heijnsbergen, D., et al. 2000, *Science* 288, 313
- Webster, A., 1995, *MNRAS*277, 1555